





Search PubMed Clear ▼ for arginine AND Xa protease AND inhibitor AND (spe Details Display Summary Save Text Clip Add Order Show: |20 Items 1-19 of 19 One page. ☐ 1: Manithody C, Yang L, Rezaie AR. Act of participal proces Role of basic residues of the autolysis loop in the catalytic function of factor Xa. Biochemistry, 2002 May 28;41(21):6780-8. PMID: 12022882 [PubMed - indexed for MEDLINE] Harry Bridge Barrell Garage Contract 厂 2: Biotin derivatives of D-Phe-Pro-Arg-CH2Cl for active-site-specific labeling of thrombin and other serine proteinases. Anal Biochem. 2001 Sep 15;296(2):254-61. PMID 11554721 [PubMed - indexed for MEDLINE] ☐ 3:00 (200 YES) (200 3:10 (a NM, 3) (3 NO, 00 a 0 NO Related Articles, Links The antithrombin P1 residue is important for target proteinase specificity but not for heparin activation of the serpin. Characterization of P1 antithrombin variants with altered proteinase specificity but normal heparin activation. Biochemistry. 2001 Jun 5;40(22):6670-9. PMID: 11380262 [PubMed - indexed for MEDLINE] 4: Turner PC, Baquero MT, Yuan S, Thoennes SR, Moyer RW. Reliated Articles, Links The cowpox virus serpin SPI-3 complexes with and inhibits urokinase-type and tissue-type plasminogen activators and plasmin. Virology. 2000 Jul 5;272(2):267-80. PMID 10873770 [PubMed - indexed for MEDLINE] **5:**St Charles R. Padmanabhan K. Arni RV, Padmanabhan KP, Tulinsk√ A. Structure of tick anticoagulant peptide at 1.6 A resolution complexed with bovine pancreatic trypsin inhibitor. Protein Sci. 2000 Feb;9(2):265-72. PMID. 10716178 [PubMed - indexed for MEDLINE] 6: Narasimhan LS, Rubin JR, Holland DR, Plummer JS, Rapundalo ST, Edmunds and programming the cost towards Jb, St-Denis Y, Siddiqui MA, Humblet C. Structural basis of the thrombin selectivity of a ligand that contains the constrained arginine mimic (2S)-2-amino-(3S)-3-(1-carbamimidoyl- piperidin-3-yl)-propanoic acid at P1. J Med Chem. 2000 Feb 10;43(3):361-8. PMID: 10669563 [PubMed - indexed for MEDLINE] O. K. St. Company Services Design, synthesis and structure-activity relationship of a series of arginine aldehyde factor Xa inhibitors. Part 1: structures based on the (D)-Arg-Gly-Arg tripeptide sequence. Bioorg Med Chem Lett. 2000 Jan 3:10(1):13-6. PMID: 10636232 [PubMed - indexed for MEDLINE]

and modelling of the bdellastasin-microplasmin system. J Mol Biol. 1999 Oct 15;293(1):93-106. PMID: 10512718 [PubMed - indexed for MEDLINE] The arrest was to ear **9:** Due MZ, Le Bonniec BF, Hopkins PC, Hipler K, Stone SR. Role of the P2 residue in determining the specificity of serpins. Biochemistry, 1996 Sep 3;35(35):11461-9. PMID 8784202 [PubMed - indexed for MEDLINE] □ 10:Lee WH, Zhang Y, Wang WY, Xiong YL, Gao R. Reductions Lies Isolation and properties of a blood coagulation factor X activator from the venom of king cobra (Ophiophagus hannah). Toxicon. 1995 Oct;33(10):1263-76. PMID: 8599178 [PubMed - indexed for MEDLINE] Exercised Articles | Lores 11:Okayama T, Seki S, Ito H, Takeshima T, Hagiwara M, Morikawa T. Lactam-conformationally restricted analogs of N alpha-arylsulfonyl arginine amide: design, synthesis and inhibitory activity toward thrombin and related enzymes. Chem Pharm Bull (Tokyo). 1995 Oct;43(10):1683-91. PMID. 8536341 [PubMed - indexed for MEDLINE] 12: Deadman JJ, Elgendy S, Goodwin CA, Green D, Baban JA, Patel G, Herate than to a set one Skordalakes E. Chino N. Claeson G. Kakkar VV. et al. Characterization of a class of peptide boronates with neutral P1 side chains as highly selective inhibitors of thrombin. J Med Chem. 1995 Apr 28;38(9):1511-22. PMID: 7739010 [PubMed - indexed for MEDLINE] 13. Padmanabhan K. Padmanabhan KP. Tulinsky A. Park CH, Bode W. Huber R. Record Art 2005. 1 or a Blankenship DT, Cardin AD, Kisiel W. Structure of human des(1-45) factor Xa at 2.2 A resolution. J Mol Biol. 1993 Aug 5;232(3):947-66. PMID: 8355279 [PubMed - indexed for MEDLINE] 14. Shuman RT, Rothenberger RB, Campbell CS, Smith GF, Gifford-Moore DS, Relating Association and the statement of the statem Gesellehen PD. Highly selective tripeptide thrombin inhibitors. J Med Chem. 1993 Feb 5;36(3):314-9 PMID: 8426361 [PubMed - indexed for MEDLINE] ☐ 15:Keyt BA, Berleau LT, Nguyen HV, Bennett WF. Re-reg Articles 1 has Radioiodination of the active site of tissue plasminogen activator: a method for radiolabeling serine proteases with tyrosylprolylarginyl chloromethyl ketone. Anal Biochem. 1992 Oct;206(1):73-83. PMID: 1456445 [PubMed - indexed for MEDLINE] 16: Dunwiddie C. Thornberry NA, Bull HG, Sardana M, Friedman PA, Jacobs Biography of the great street and the control of the c JW, Simpson F Antistasin, a leech-derived inhibitor of factor Xa. Kinetic analysis of enzyme inhibition and identification of the reactive site. J Biol Chem. 1989 Oct 5;264(28):16694-9. PMID: 2777803 [PubMed - indexed for MEDLINE] T 17: Powers JC, Kam CM, Narasimhan L, Oleksyszyn J, Hernandez MA, Ueda T, For the additional trans-Mechanism-based isocoumarin inhibitors for serine proteases: use of active site structure and substrate specificity in inhibitor design. J Cell Biochem. 1989 Jan; 39(1):33-46. Review. PMID: 2654146 [PubMed - indexed for MEDLINE]

Structure of the complex of the antistasin-type inhibitor bdellastasin with trypsin

	ura R. Robison DJ. B	•			Harangan, a	•
	1 ,	nethanesulton	yl fluoride.	an irrever	sible inhibitor of serine	
	oteases.	11.10.31. 1050	. 1			
	ochemistry, 1980 Oct					
PN	IID: 6775696 [PubMc	ed - indexed for i	MEDLINE			
□ 19: H;	jikata A. Okamoto S.	Kikumoto R. La	mao Y.		An energy are as a	
Ki	netic studies on th	e selectivity o	of a synthet:	ie thrombi	n-inhibitor using syntheti	.c
pe	ptide substrates.					
•	romb Haemost. 1979	Oct 31;42(3):10.	39-45.			
PN	IID: 159513 [PubMed	l - indexed for N	(EDLINE)			
	•		•			
<u> </u>	l o			11		
Display	Summary	Sort	▼ Save	lext	Clip Add Order	

Write to the Help Desk NCBI | NLM | NIH Department of Health & Human Services Freedom of Information Act | Disclaimer One page.

Items 1-19 of 19

Show: 20 ▼







Search PubMed	▼ for arginine AND Xa protease AND inhibitor AND sper Go Clear							
	Commission of Assistance of the Commission of Commission o							
	Display Summary Sort Save Text Clip Add Order							
	Show: 20 Items 1-15 of 15 One page							
	T 1: Manthody C. Yang L. Rezaie AR. Role of basic residues of the autolysis loop in the catalytic function of factor Xa. Biochemistry. 2002 May 28:41(21):6780-8. PMID 12022882 [PubMed - indexed for MEDLINE]							
	□ 2: Volume of the transfer of the Control of the							
	Biotin derivatives of D-Phe-Pro-Arg-CH2Cl for active-site-specific labeling of thrombin and other serine proteinases. Anal Biochem. 2001 Sep 15;296(2):254-61. PMID 11554721 [PubMed - indexed for MEDLINE]							
	□ 3:0 June AMS Justic ReB = SV, 0 Just Sc. O vo. St. Because Andres, Line							
Rough Rough And Comment of the Comme	The antithrombin P1 residue is important for target proteinase specificity but not for heparin activation of the serpin. Characterization of P1 antithrombin variants with altered proteinase specificity but normal heparin activation. Biochemistry. 2001 Jun 5;40(22):6670-9. PMID 11380262 [PubMed - indexed for MEDLINE]							
	The cowpox virus serpin SPI-3 complexes with and inhibits urokinase-type and tissue-type plasminogen activators and plasmin. Virology. 2000 Jul 5:272(2):267-80. PMID. 10873770 [PubMed - indexed for MEDLINE]							
	Structure of tick anticoagulant peptide at 1.6 A resolution complexed with bovine pancreatic trypsin inhibitor. Protein Sci. 2000 Feb;9(2):265-72. PMID. 10716178 [PubMed - indexed for MEDLINE]							
	Design, synthesis and structure-activity relationship of a series of arginine aldehyde factor Xa inhibitors. Part 1: structures based on the (D)-Arg-Gly-Arg tripeptide sequence. Bioorg Med Chem Lett. 2000 Jan 3:10(1):13-6. PMID 10636232 [PubMed - indexed for MEDLINE]							
	7:Rester U. Bode W. Moser M. Parry MA. Huber R. Auerswaid F. Green and M. Green and M. Structure of the complex of the antistasin-type inhibitor bdellastasin with trypsin and modelling of the bdellastasin-microplasmin system. J. Mol Biol. 1999 Oct 15:293(1):93-106. PMID: 10512718 [PubMed - indexed for MEDLINE]							
	■ 8: Dire MZ, Le Bonniec BF, Hopkins PC, Hipler K, Stone SR.							

Role of the P2 residue in determining the specificity of serpins.

Biochemistry, 1996 Sep 3;35(35):11461-9.

9:Lee WH, Zhang Y, Wang WY, Xiong YL, Gao R.	And great articles of the			
Isolation and properties of a blood coagulation factor X activator	r from the venom of			
king cobra (Ophiophagus hannah).				
Toxicon, 1995 Oct;33(10):1263-76. PMID: 8599178 [PubMed - indexed for MEDLINE]				
,				
10: Deadman JJ, Flgendy S, Goodwin CA, Green D, Baban JA, Patel G, Skordalakes E, Chino N, Claeson G, Kakkar VV, et al.	An object the party of the control o			
Characterization of a class of peptide boronates with neutral P1	side chains as			
highly selective inhibitors of thrombin. J Med Chem. 1995 Apr 28:38(9):1511-22.				
PMID 7739010 [PubMed - indexed for MEDLINE]				
The state of the s				
11. Padmanabhan K, Padmanabhan KP, Tulinsky A, Park CH, Bode W, Huber R Blankenship DT, Cardin AD, Kisiel W.	er nye late it Arthoryes (Elifeks). L			
Structure of human des(1-45) factor Xa at 2.2 A resolution.				
J Mol Biol. 1993 Aug 5;232(3):947-66. PMID: 8355279 [PubMed - indexed for MEDLINE]				
1. Mile 1030027% (Fub. Med Indexed for Med Eliste)				
☐ 12:Keyt BA, Berleau LT, Nguyen HV, Bennett WF.	Related Articles, Lines			
Radioiodination of the active site of tissue plasminogen activato				
radiolabeling serine proteases with tyrosylprolylarginyl chlorom	ethyl ketone.			
Anal Biochem. 1992 Oct;206(1):73-83. PMID: 1456445 [PubMed - indexed for MEDLINE]				
13: Dunwiddie C, Thornberry NA, Bull HG, Sardana M, Friedman PA, Jacobs JW, Simpson E.	The standard and Errors			
Antistasin, a leech-derived inhibitor of factor Xa. Kinetic analys	is of enzyme			
inhibition and identification of the reactive site. J Biol Chem. 1989 Oct 5;264(28):16694-9.				
PMID: 2777803 [PubMed - indexed for MEDLINE]				
14:Powers JC, Kam CM, Narasimhan L, Oleksyszyn J, Hernandez MA, Ueda T				
Mechanism-based isocoumarin inhibitors for serine proteases: u	se of active site			
structure and substrate specificity in inhibitor design. J Cell Biochem. 1989 Jan;39(1):33-46 Review.				
PMID 2654146 [PubMed - indexed for MEDLINE]				
T 15-Loren D. Dobieses INI. Ding INI.	Company to a second control			
(p-Amidinophenyl)methanesulfonyl fluoride, an irreversible inh	- Related Articles Links ibitor of serine			
proteases.	ionor or serme			
Biochemistry, 1980 Oct 14;19(21):4859-64.				
PMID 6775696 [PubMed - indexed for MEDLINE]				
Display Summary ▼ Sort ▼ Save Text Clip Ac	dd Order			
Show. 20 <u>▼</u> Items 1-15 of 15	One page.			

Write to the Help Desk NCBI | NLM | NIH Department of Health & Human Services Freedom of Information Act | Disclaimer







Search PubMed
▼ for arginine AND Xa protease AND inhibitor AND (spe Preview Go Clear History

• Search History will be lost after one hour of inactivity.

• To combine searches use # before search number, e.g., #2 AND #6.

• Search numbers may not be continuous; all searches are represented.

Searc	Most Recent Queries	Time	Result
#	4 Search arginine AND Xa protease AND inhibitor AND (specificity or selectivity)	10:46:53	19
#	3 Search arginine AND Xa protease AND inhibitor AND specificity	10:44:44	15
#	2 Search arginine AND Xa protease AND specificity	10:44:38	35
#	1 Search	10:43:59	()

Clear History

Write to the Help Desk NCBI | NEM | NIH Department of Health & Human Services Freedom of Information Act | Disclaimer